

## Cholinesterase Inhibitor Therapy for Alzheimer's Disease Criteria for Use

VHA Pharmacy Benefits Management Strategic Healthcare Group and the Medical Advisory Panel

*The following recommendations are dynamic and will be revised, as new clinical data become available. These criteria are not intended to interfere with clinical judgment. Rather, they are intended to assist practitioners in providing cost effective, consistent, high quality care.*

Alzheimer's disease (AD) is the most common cause of dementia in North America. The first clinical signs of AD are impairments of memory, language and visuospatial function, some of which can be explained by loss of cholinergic neurons in the basal forebrain. This loss contributes to the symptom development of AD. The clinical diagnosis of AD can be made using the criteria of the Diagnostic and Statistical Manual – IV (American Psychiatric Association, 1994)<sup>40</sup>. Once the diagnosis of AD has been made, the severity of the dementia can be estimated and monitored using a cognitive instrument such as the Mini-Mental State Examination (MMSE).<sup>39</sup> The main pharmacological approach to limiting cognitive and functional decline in AD is to increase synaptic levels of acetylcholine through use of cholinesterase inhibitors (CIs). Treatment of AD should take into account concerns about quality of life for the patient and for members of his/her family. Therefore, it is important to respect patient and family preferences when making treatment recommendations. Since these agents are costly and are sometimes associated with side effects, their role in AD management must be clearly defined. The use of CI in other disease states such as Lewy Body and vascular dementia is not addressed by this document.

**The CIs, donepezil (Aricept®), galantamine (Reminyl®) and rivastigmine (Exelon®) are modestly effective in delaying cognitive and functional decline in patients with mild to moderate AD.**<sup>1-4, 7-21</sup>

There is no level I evidence\* to demonstrate superiority amongst the agents. Dose and titration schedules differ among the agents (see Table 1). Generally the agents are well tolerated with common adverse effects managed with titration and dose adjustments. CIs have been shown to offer benefit by delaying nursing home placement or by reducing costs of care in the home.<sup>22-36</sup> In one published trial, donepezil has also been shown to delay functional decline in patients with moderate to severe AD who are still living in the community.<sup>12</sup> Data on the effects of CI use in nursing home patients with mild to moderate disease are not available. Therefore, the initiation of CI in the nursing home population must consider the cost effectiveness of these agents and paucity of data on their benefit in this group. Tacrine (Cognex®) is no longer used as a first-line agent in AD because of the risk of hepatotoxicity. There is insufficient evidence\* to support the initiation of CIs in AD patients with severe dementia (MMSE<10). **CI therapy should be discontinued if there is evidence of a) poor compliance; b) persistent side effects; c) mutual agreement between caregiver and provider. A tapered discontinuation is recommended in a) progression to severe dementia (MMSE<10); b) development of a concomitant contraindicated medical condition (see algorithm); c) permanent loss of caregiver or d) permanent institutionalization for severe dementia with functional loss.** CI therapy may be re-initiated in patients where there is evidence of a rapid decline upon discontinuation of CI therapy. There is insufficient evidence\* to support switching between CI's if one agent is not tolerated or is ineffective.<sup>6</sup>

**Table 1: Cholinesterase Inhibitors#**

Drug	Initial dose	Titration schedule every	Recommended dose	Cost/month	Minimum Therapeutic Dose	Formulations
<b>Donepezil**</b> Aricept®	5 mg QD	4 weeks	10 mg QD	\$76.50	5 mg QD	Tablets: 5,10 mg
<b>Galantamine</b> Reminyl®	4 mg BID	4 weeks	12 mg BID	\$78.00	8 mg BID	Tablets: 4,8,12 mg Oral liquid 4mg/ml
<b>Rivastigmine</b> Exelon®	1.5mg BID	4 weeks	6mg BID	\$80.40	3 mg BID	Capsules: 1.5,3,4.5, 6 mg Oral liquid 2mg/ml

#Cognex® (tacrine) is not included in this table due to hepatotoxicity risk

\*\* Currently only agent on VA National Formulary

QD- once daily, BID- twice daily

### \* Quality of Evidence

**I:** Evidence obtained from at least one properly randomized controlled trial. US Preventative Task Force scale Am J Prev Med 2002;20(3S):21-35.

# Algorithm for CI Therapy in Alzheimer's Dementia




Table 2: Anticholinergic Activity of Selected Agents<sup>@</sup>

Class	none- minimal anticholinergic effect	moderate anticholinergic effect	high anticholinergic effects
<b>Antidepressants</b>	Bupropion, sertraline, paroxetine citalopram, fluoxetine, fluvoxamine	nortriptyline, desipramine , doxepin, maprotiline mirtazapine	amitriptyline, protriptyline, clomipramine, amoxapine, imipramine,
<b>Antiparkinsonian</b>	levodopa, pergolide, pramipexole, ropinirole	amantadine	benztropine, trihexyphenidyl
<b>Antipsychotics</b>	risperidone ziprasidone haloperidol	olanzapine, quetiapine, chlorpromazine, pimozide	clozapine, thioridazine, mesoridazine, promazine, triflupromazine
<b>Antispasmodics</b>		belladonna alkaloids, dicyclomine tolterodine, diphenoxylate	atropine, oxybutynin
<b>Antihistamines</b>	loratadine, fexofenadine, cetirizine	brompheniramine, chlorpheniramine hydroxyzine	clemastine, diphenhydramine, promethazine
<b>Other</b>			meclizine, scopolamine, dimenhydrat, prochlorperazine

<sup>@</sup>It should be remembered that patients with AD are uniquely sensitive to anticholinergic agents and even minimal effects can translate into clinical changes. These agents should be avoided if at all possible.

Table 3: Drug Interactions

Precipitant drug	Object drug	Description
Cimetidine	Galantamine	Bioavailability of galantamine increased by 16%*
Ketoconazole	Galantamine	Galantamine AUC increased 30%*
Paroxetine	Galantamine	Galantamine oral bioavailability increased 40%*
Erythromycin	Galantamine	Galantamine AUC increased 10%*
Donepezil	Furosemide, digoxin, warfarin	Donepezil did not alter the binding of the object drugs
Ketoconazole	Donepezil	Donepezil metabolism may be altered#
Quinidine	Donepezil	Donepezil metabolism may be altered#
CI class	Succinylcholine	Exaggeration of relaxation response

*Donepezil and Galantamine* are metabolized through both CYP 450 3A4 and 2D6 isoenzymes

*Rivastigmine* is not extensively metabolized by the cytochrome P450 system

# Clinical effects of the interactions are unknown; they are postulated interactions.

\* Clinical significance of these interactions is unknown.

AUC-Area Under the Concentration Time Curve

## References

1. Doody RS, Stevens JC, Beck C, Dubinsky RM, Kaye JA, et al. Practice Parameter: Management of Dementia (an evidence based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2001;56:1154-1166.
2. Rogers SL, Doody Rs, Pratt RD, Ieni JR. Long-term efficacy and safety of donepezil in the treatment of AD: a final analysis of a US multicenter open-label study. *European Neuropsychopharmacology* 2000 May;10(3): 195-203.
3. Homma A, Takeda M, Imai Y, Ukada F, Hasegawa K, et al. Clinical efficacy and safety of donepezil on cognitive and global function in patients with AD. A 24 week, multicenter, double-blind, placebo-controlled study in Japan. E2020 group. *Dementia & Geriatric Cognitive Disorders* 2000 Nov-Dec;11(6):299-313.
4. Farlow MR. A 52-week study of the efficacy of rivastigmine in patients with mild to moderately severe AD *European Neurology* 2000;44(4):236-41.
5. Forette F, Anand R, Gharabawi G. A phase II study in patients with AD to assess the preliminary efficacy and maximum tolerated dose of rivastigmine. *European Journal of Neurology* 1999 Jul;6(4):423-9.
6. Auriacombe S, Pere JJ, Loria-Kanza Y, Vellas B. Efficacy and Safety of Rivastigmine in Patients with Alzheimer's who failed to benefit from Treatment with Donepezil. *Current Medical Research and Opinion* 2002;18(3):129-138.
7. Wilkinson DG, Passmore AP, Bullock R et al. A Multinational, Randomized, 12 week, Comparative Study of Donepezil and Rivastigmine in Patients with Mild to Moderate Alzheimer's Disease. *Int J Clin Pract*;56(6):441-446
8. Burns A, Rossor M, Hacker J, Gauthier S, Petit H, et al. The effects of donepezil in Alzheimer's disease-results from a multinational trial. *Dementia & Geriatric Cognitive Disorders* 1999 May-Jun;10(3):237-44.
9. Feldman H, Gauthier S, Hecker J, Vellas B, Subbiah P, et al. A 24 week, randomized, double-blind study of donepezil in moderate to severe AD. *Neurology* 2001 Aug 28;57(4):613-20.
10. Doody RS, Dunn JK, Clark CM, Farlow M, Foster NL, et al. Chronic donepezil treatment is associated with slowed cognitive decline in AD. *Dementia & Geriatric Cognitive Disorders* 2001 Jul-Aug;12(4):295-300.
11. Doody RS, Geldmacher DS, Gordon B, Perdomos CA, Pratt RP ,et al. Open-label, multicenter, phase 3 extension study of the safety and efficacy of donepezil in patients with AD. *Archives of Neurology*. 2001 Mar;58(3):427-33.
12. Tariot PN, Cummings JL, Katz IR, Mintzer J, Perdomo CA, et al.. A randomized double-blind, placebo-controlled study of the efficacy and safety of donepezil. *J Am Geriatr Soc* 2001 Dec;49(12):1590-9.
13. Winblad B, Engedal K, Soininen H, Verhay F, Waldemar G, et al. A 1- yr randomized placebo-controlled study of donepezil in patients with mild to moderate AD. *Neurology* 2001 Aug 14;57(3):489-95.
14. Farlow MR, Hake A, Messina J, Hartman R, Veach J, et al. Response of patients with AD to rivastigmine treatment is predicted by the rate of disease progression. *Archives of Neurology* 2001 Mar;58(3):417-22.
15. Rosler M, Anand R, Cicin-Sain A, Gauthier S, Agid Y, Dal-Bianco P, Stahelin HB, Hartman R, Gharabawi M. Efficacy and safety of rivastigmine in patients with Alzheimer's disease: international randomised controlled trial. *BMJ*. 1999 Mar 6;318(7184):633-8.

16. Wilkinson D, Murray J. Galantamine: a randomized, double-blind, dose comparison in patients with AD. *International Journal of Geriatric psychiatry*. 2001 Sep;16(9):852-7.
17. Wilcock GK, Lilienfeld S, Gaens E. Efficacy and safety of galantamine in patients with mild to moderate Alzheimer's disease: multicentre randomized controlled trial. Galantamine International-1 Study Group. *BMJ* 2000 Dec 9;321(7274):1445-9.
18. Tariot PN, Solomon PR, Morris JL, Kershaw P, Lilienfeld S, et al. A 5-month, randomized, placebo-controlled trial of galantamine in AD. The Galantamine USA-10 Study Group. *Neurology* 2000 Jun 27;2269-76.
19. Raskind MA, Peskind ER, Wessel T, Yuan W. Galantamine in AD: A 6-month randomized, placebo-controlled trial with a 6-month extension. The Galantamine USA-1 Study Group. *Neurology* 2000 Jun 27;54(12):2261-8.
20. Rockwood K, Mintzer J, Truyen L, Wessel T, Wilkinson D. Effects of a flexible galantamine dose in Alzheimer's disease: a randomized, controlled trial. *J Neurol Neurosurg Psychiatry*. 2001 Nov;71(5):589-95.
21. Erkinjuntti T, Kurz A, Gauthier S, Bullock R, Lilienfeld S, Damaraju CV. Efficacy of galantamine in probable vascular dementia and Alzheimer's disease combined with cerebrovascular disease: a randomized trial. *Lancet*. 2002 Apr 13;359(9314):1283-90.
22. Stewart A, Phillips R, Dempsey G. Pharmacotherapy for people with Alzheimer's disease: a Markov-cycle evaluation of five years' therapy using donepezil. *Int J Geriatr Psychiatry*. 1998 Jul;13(7):445-53.
23. Small GW, Donohue JA, Brooks RI. An economic evaluation of donepezil in the treatment of Alzheimer's disease. *Clin Ther*. 1998 Jul-Aug;20(4):838-50.
24. Jonsson L, Lindgren P, Wimo A, Jonsson B, Winblad B. The cost-effectiveness of donepezil therapy in Swedish patients with Alzheimer's disease: a Markov model. *Clin Ther*. 1999 Jul;21(7):1230-40.
25. O'Brien BJ, Goeree R, Hux M, Iskedjian M, Blackhouse G, et al. Economic evaluation of donepezil for the treatment of Alzheimer's disease in Canada. *J Am Geriatr Soc*. 1999 May;47(5):570-8.
26. Neumann PJ, Hermann RC, Kuntz KM, Araki SS, Duff SB, et al. Cost-effectiveness of donepezil in the treatment of mild or moderate Alzheimer's disease. *Neurology*. 1999 Apr 12;52(6):1138-45.
27. Fillit H, Guterman EM, Lewis B. Donepezil use in managed Medicare: effect on health care costs and utilization. *Clin Ther*. 1999 Dec;21(12):2173-85.
28. Hill JW, Futerman R, Mastey V, Fillit H. The effect of donepezil therapy on health costs in a Medicare managed care plan. *Manag Care Interface*. 2002 Mar;15(3):63-70.
29. Ikeda S, Yamada Y, Ikegami N. Economic evaluation of donepezil treatment for Alzheimer's disease in Japan. *Dement Geriatr Cogn Disord*. 2002;13(1):33-9.
30. Fenn P, Gray A. Estimating long-term cost savings from treatment of Alzheimer's disease. A modelling approach. *Pharmacoeconomics*. 1999 Aug;16(2):165-74.
31. Hauber AB, Gnanasakthy A, Snyder EH, Bala MV, Richter A, et al. Potential savings in the cost of caring for Alzheimer's disease. Treatment with rivastigmine. *Pharmacoeconomics*. 2000 Apr;17(4):351-60.

32. Baladi JF, Bailey PA, Black S, Bouchard RW, et al. Rivastigmine for Alzheimer's disease: Canadian interpretation of intermediate outcome measures and cost implications. *Clin Ther*. 2000 Dec;22(12):1549-61.
33. Hauber AB, Gnanasakthy A, Mauskopf JA. Savings in the cost of caring for patients with Alzheimer's disease in Canada: an analysis of treatment with rivastigmine. *Clin Ther*. 2000 Apr;22(4):439-51.
34. Getsios D, Caro JJ, Caro G, Ishak K. Assessment of health economics in Alzheimer's disease (AHEAD): galantamine treatment in Canada. *Neurology*. 2001 Sep 25;57(6):972-8.
35. Garfield FB, Getsios D, Caro JJ, Wimo A, Winblad B. Assessment of Health Economics in Alzheimer's Disease (AHEAD): treatment with galantamine in Sweden. *Pharmacoeconomics*. 2002;20(9):629-37.
36. Caro JJ, Slas M, Ward A, Getsios D, Mehnert A. Economic analysis of galantamine, a cholinesterase inhibitor, in the treatment of patients with mild to moderate Alzheimer's disease in the Netherlands. *Dement Geriatr Cogn Disord*. 2002;14(2):84-9.
37. Gray SL, Lai KV, Larson EB. Drug-induced cognition disorders in the elderly. *Drug Safety* 1999 Aug 21 (20):101-122.
38. Leung GM, Yeung RYT, Chi I, Chu LW. The economics of Alzheimer's disease. *Dement Geriatr Cogn Disord* 2003 Jan; 15:34-43.
39. Folstein NF, Folstein SE, McHugh PR. Mini-mental State: a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189-198.
40. American Psychiatric Association (APA). *Diagnostic and Statistical Manual of Mental Disorders-IV*. APA, Washington, DC. 1994.

Prepared by: Kavita Palla Pharm D, Kathryn Tortorice Pharm D, BCPS

Reviewed by: Linda Hershey M.D., William Korchik M.D., J. Riley McCarten M.D.,  
Susan Cooley PhD., Kye Kim M.D, Ladislav Volicer, M.D., PhD